

Citation for published version:

Bulla, M, Valcu, M, Dokter, AM, Dondua, AG, Kosztolányi, A, Rutten, AL, Helm, B, Sandercock, BK, Casler, B, Ens, BJ, Spiegel, CS, Hassell, CJ, Küpper, C, Minton, C, Burgas, D, Lank, DB, Payer, DC, Loktionov, EY, Nol, E, Kwon, E, Smith, F, Gates, HR, Vitnerová, H, Prüter, H, Johnson, JA, St Clair, JJH, Lamarre, JF, Rausch, J, Reneerkens, J, Conklin, JR, Burger, J, Liebezeit, J, Bêty, J, Coleman, JT, Figuerola, J, Hooijmeijer, JCEW, Alves, JA, Smith, JAM, Weidinger, K, Koivula, K, Gosbell, K, Exo, KM, Niles, L, Koloski, L, McKinnon, L, Praus, L, Klaassen, M, Giroux, MA, Sládecek, M, Boldenow, ML, Goldstein, MI, Sálek, M, Senner, N, Rönkä, N, Lecomte, N, Gilg, O, Vincze, O, Johnson, OW, Smith, PA, Woodard, PF, Tomkovich, PS, Battley, PF, Bentzen, R, Lanctot, RB, Porter, R, Saalfeld, ST, Freeman, S, Brown, SC, Yezerinac, S, Székely, T, Montalvo, T, Piersma, T, Loverti, V, Pakanen, VM, Tijssen, W & Kempenaers, B 2016, 'Unexpected diversity in socially synchronized rhythms of shorebirds', *Nature*, vol. 540, no. 7631, pp. 109-113.
<https://doi.org/10.1038/nature20563>

DOI:

[10.1038/nature20563](https://doi.org/10.1038/nature20563)

Publication date:

2016

Document Version

Peer reviewed version

[Link to publication](#)

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

1 **Defying the 24-h day: Unexpected diversity in socially synchronized rhythms of shorebirds**

2 Martin Bulla¹, Mihai Valcu¹, Adriaan M. Dokter, Alexei G. Dondua, András Kosztolányi, Anne Rutten¹, Barbara Helm, Brett K. Sandercock,
3 Bruce Casler, Bruno J. Ens, Caleb S. Spiegel, Chris J. Hassell, Clemens Küpper, Clive Minton, Daniel Burgas, David B. Lank, David C. Payer,
4 Egor Y. Loktionov, Erica Nol, Eunbi Kwon, Fletcher Smith, H. River Gates, Hana Vítnerová, Hanna Prüter, James A. Johnson, James J. H. St
5 Clair, Jean-François Lamarre, Jennie Rausch, Jeroen Reneerkens, Jesse R. Conklin, Joanna Burger, Joe Liebezeit, Joël Bêty, Jonathan T.
6 Coleman, Jordi Figuerola, Jos C. E. W. Hooijmeijer, José A. Alves, Joseph A. M. Smith, Karel Weidinger, Kari Koivula, Ken Gosbell, Larry
7 Niles, Laura Koloski, Laura McKinnon, Libor Praus, Marcel Klaassen, Marie-Andrée Giroux, Martin Sládeček, Megan L. Boldenow, Michael
8 Exo, Michael I. Goldstein, Miroslav Šálek, Nathan Senner, Nelli Rönkä, Nicolas Lecomte, Olivier Gilg, Orsolya Vincze, Oscar W. Johnson,
9 Paul A. Smith, Paul F. Woodard, Pavel S. Tomkovich, Phil Battley, Rebecca Bentzen, Richard B. Lanctot, Ron Porter, Sarah T. Saalfeld, Scott
10 Freeman, Stephen C. Brown, Stephen Yezerinac, Tamás Székely, Theunis Piersma, Tomás Montalvo, Vanessa Loverti, Veli-Matti Pakanen,
11 Wim Tijssen, Bart Kempenaers¹

12 **All organisms exhibit alternating phases of different behaviours. These biological rhythms are thought to be**
13 **under strong selection, influenced by the rhythmicity of the environment. Behavioural rhythms are well**
14 **studied in isolated individuals under laboratory conditions^{1,2}, but in free-living populations, individuals have**
15 **to temporally synchronize their activities with those of others. The behavioural rhythms that emerge from**
16 **such social synchronization and the underlying evolutionary and ecological drivers that shape them are**
17 **poorly understood²⁻⁵. Here, we address this for a particularly sensitive phase of social synchronization when**
18 **pair members care for their offspring⁶, and thereby potentially compromise their individual rhythms.**
19 **Specifically, we use data from 729 nests of 91 populations of 32 biparentally-incubating shorebird species,**
20 **where parents synchronize to achieve continuous coverage of developing eggs. We found remarkable within-**
21 **and between-species diversity in incubation rhythms. Between species, the median length of one parent's**
22 **incubation bout varied from one to 19 hours, while period length – the cycle of female and male probability**
23 **to incubate – varied from six to 43 hours. The length of incubation bouts was unrelated to variables reflecting**
24 **energetic demands, but species relying on crypsis had longer incubation bouts than those that are readily**
25 **visible or actively protect their nest against predators. Rhythms strictly entrainable to the 24-h light-dark**

¹Department of Behavioural Ecology and Evolutionary Genetics, Max Planck Institute for Ornithology, Eberhard Gwinner Str, 82319 Seewiesen, Germany.
Affiliations of the other authors are given in the Supplementary Data 1.

cycle were less likely at high latitudes and absent in 18 species. Our results indicate that even under similar environmental conditions and despite 24-h environmental cues, social synchronization can generate far more diverse behavioural rhythms than expected from studies of individuals in captivity. The risk of predation, not the risk of starvation, may be a key factor underlying the diversity in these rhythms.

The activity rhythms of organisms, from bacteria to humans, are typically the output of internal ‘biological clocks’ synchronized to the daily environmental cycle¹. These biological clocks and the resulting rhythms are largely genetically determined and conserved across species⁷⁻⁹. Yet, behavioural rhythms of individuals are further shaped by the social environment, including potential mates, allies, competitors, prey, or predators^{3-5,10,11}. As a result of such social interactions, individuals can temporally segregate their daily activities (e.g. prey avoiding predators, subordinates avoiding dominants) or synchronize their activities (e.g. foraging in group, communal defence against predators, pairs reproducing or caring for offspring)^{3-5,12,13}. Although social synchronization of conspecifics has been documented across taxonomic groups²⁻⁵ and although the behavioural rhythms that emerge from social synchronization have occasionally been studied in captive individuals^{2-5,14}, we know little about socially emerged rhythms in nature^{3-5,10,11}, and even less about the factors that influence them such as evolutionary history, energetic demands, predation risk and periodicity in the environment.

Here, we report on a large-scale study of socially emerged rhythms and their correlates in the wild, using biparental incubation in shorebirds as a model. Incubation by both parents prevails in almost 80% of non-passerine families¹⁵ and is the most common form of care in shorebirds¹⁶. Biparental shorebirds are typically monogamous¹⁷, mostly lay three or four eggs in an open nest on the ground¹⁷, and cover their eggs almost continuously¹⁵. Pairs achieve this through synchronization of their activities such that one of them is responsible for the nest at a given time (i.e. an incubation bout). Alternating female and male bouts generate an incubation rhythm with a specific period length (cycle of high and low probability for a parent to incubate).

We collected data on incubation rhythms from 91 populations of 32 shorebird species belonging to 10 genera (Fig. 1), extracted the length of 34,225 incubation bouts from 729 nests, and determined the period length for pairs in 584 nests (see Extended Data Fig. 1).

We found vast between- and within- species variation in incubation bout length and in period length (Fig. 2-4 & Extended Data Fig. 2). Different species, but also different pairs of the same species, adopted strikingly different incubation rhythms, even when breeding in the same area (see, e.g. incubation rhythms in Barrow, Alaska, represented by ① in Fig. 2; incubation rhythms for each nest are in Supplementary Actograms¹⁸). Whereas in some pairs parents exchanged incubation duties about 20 times a day (Fig. 3a; e.g. *Charadrius semipalmatus*, Fig. 2a), in others a single parent regularly incubated for 24 hours (Fig. 3a; e.g. *Limnodromus scolopaceus*, Fig. 2a), with exceptional bouts of up to 50 hours (Supplementary Actograms¹⁸). Similarly, whereas incubation rhythms of pairs in 22% of nests followed a strict 24-h period (Fig. 3b; e.g. *Tringa flavipes*, Fig. 2a), the rhythms of others dramatically deviated from a 24-h period (Fig. 3b) resulting in ultradian (<20-h in 12% of nests; e.g. *Numenius phaeopus*; Fig. 2a), free-running like (e.g. *Calidris alpina*; Fig. 2a) and infradian rhythms (>28-h in 8% of nests), some with period lengths up to 48-h (e.g. *Limnodromus scolopaceus*; Fig. 2a). This variation in period length partly related to the variation in bout length (Fig. 4): in the suborder Scolopaci period length correlated positively with median bout length, but in the suborder Charadrii species with 24-h periods had various bout lengths, and species with similar bout lengths had different period lengths (Fig. 4).

Despite substantial within-species variation, we found a strong evolutionary signal for both bout and period length with a coefficient of phylogenetic signal λ close to 1 (Extended Data Table 1). This is consistent with the notion that biological rhythms are largely genetically determined and conserved among related species⁸⁻¹⁰. However, the phylogenetic effect seems unevenly distributed over taxonomic level. Suborder explained 33% of the phenotypic variance in both bout and period length, with the Scolopaci having longer incubation bouts and periods than the Charadrii (Extended Data Table 2; Fig. 3 & 4). Species explained 41% of the phenotypic variation in bout length and 46% in period length, but genus explained little (<1% in both; Extended Data Table 2), suggesting that despite a strong phylogenetic signal, these traits can rapidly diverge (Fig. 3c).

Two ecological factors may explain the observed variation in bout length. First, the ‘energetic demands hypothesis’ stipulates that the length of an incubation bout depends on a bird’s energetic state^{15,19}. This

predicts that (1) large species will have longer incubation bouts than smaller species, because they radiate less body heat per unit of mass and (2) incubation bouts will shorten with increasing breeding latitude, because – everything else being equal – energy stores will deplete faster in colder environments (Extended Data Fig. 3a-b shows latitudinal cline in summer temperatures). However, bout length was unrelated to body size (Fig. 5a) and correlated positively (instead of negatively) with latitude (Fig. 5b). These correlational results across populations and species support recent experimental findings within species²⁰ and suggest that in biparentally-incubating shorebirds energetic demands are not an important ecological driver underlying variation in bout length.

An alternative explanation for variation in the length of incubation bouts relates to anti-predation strategies. Those species that rely primarily on parental crypsis (Extended Data Fig. 4a) benefit from reduced activity near the nest, because such activity can reveal the nest's location to potential predators^{21,22}. Thus, in these species, selection will favour fewer change-overs at the nest, and hence longer incubation bouts. In contrast, species that are clearly visible when sitting on the nest or that rely on active anti-predation behaviour (Extended Data Fig. 4b), including having a partner on the watch for predators, leaving the nest long before the predator is nearby and mobbing the predator¹⁷, obtain no advantage from minimizing activity. For these species, bout length can shorten, which may be advantageous for other reasons (e.g. reduced need to store fat). We quantified anti-predation strategy as the distance at which the incubating parent left the nest when approached by a human (escape distance), because cryptic species stay on the nest longer (often until nearly stepped upon)¹⁷. Despite the large geographical distribution of the studied species, with related variability in the suite of predators and predation pressure²³, and even when controlling for phylogeny (which captures much of the variation in anti-predation strategy, Extended data Fig. 5), escape distance negatively correlated with the length of incubation bouts (Fig. 5c). This result suggests that bout length co-evolved with the anti-predation strategy.

Under natural conditions, most organisms exhibit 24-h rhythmicity, but during summer, when most shorebirds breed, the 24-h variation in light decreases with latitude leading to continuous polar daylight²⁴ (Extended Data Fig. 4c-d). Such reduced variation in 24-h light intensity may cause a loss of 24-h rhythmicity^{25,26}. As a consequence, circadian behavioural rhythms should exhibit a latitudinal cline²⁴. As predicted, incubation

rhythms with periods that do not follow the 24-h light-dark cycle, such as ‘free running-like patterns’ (left column in Fig. 2), occurred more often in shorebirds breeding at higher latitudes (Fig. 5d). The absolute deviations of periods from 24-h and 24-h harmonics also increased with latitude (Fig. 5e; Extended Data Table 3). Although this supports the existence of a latitudinal cline in socially emerged behavioural rhythms²⁴, we found a substantial number of rhythms that defy the 24-h day even at low and mid latitudes (Fig 5d-e).

Many shorebirds predominantly use tidal habitats, at least away from their breeding ground¹⁷. To anticipate tidal foraging opportunities, these species may have activity patterns with a period length resembling the tidal period. As changing to a different rhythm is costly²⁷, these tidal activity patterns might carry over to incubation. Although half of our species are tidal away from their breeding grounds, and some forage in tidal areas also during breeding (~10% of populations), in only 5% of nests did pairs display a period length that can be entrained by the tide. Moreover, tidal species had similar (not longer) periods than non-tidal ones (Extended Data Table 3). Hence, unlike the 24-h light-dark cycle, tidal life-history seems to play at best a negligible role in determining incubation rhythms.

Three main questions arise from our results. First, is variation in incubation bout length in cryptic species related to the actual predation pressure? This can be tested by comparing bout length between populations of a particular species that are exposed to different predator densities, or between years that differ in predation pressure. Second, it remains unclear how the diverse social rhythms emerge. Are these rhythms a consequence of behavioural flexibility, or a ‘fixed’ outcome of synchronization between the circadian clocks of the two individuals involved? An experimental study on ring doves (*Streptopelia risoria*) suggests that parents may even use two timers - circadian oscillation and interval timing - to determine when to incubate²⁸. Parents rapidly adjusted their schedules to phase-shifted photoperiods and their incubation rhythm ‘free-ran’ in constant dim illumination (implying a circadian mechanism), whereas an experimental delay in the onset of an incubation bout did not change the length of the bout because the incubating parent refused to leave the nest until its incubation bout reached the ‘typical’ duration (implying interval timing). Third, what – if any – are the fitness consequences for the parents of having a certain incubation rhythm? For example, the costs of having a particular incubation rhythm may be unevenly distributed between the two parents (e.g. because one parent is

on incubation duty when food is more readily available, or because one parent ‘enforces’ its own rhythm at a cost to the other parent).

In conclusion, our results reveal that under natural conditions social synchronization can generate much more diverse rhythms than expected from previous work, and that these rhythms often defy the assumptions of entrainment to the 24-h day-night cycle. Not risk of starvation, but risk of predation seems to play a key role in determining some of the variation in incubation rhythms. We describe this diversity in the context of biparental incubation, but diverse behavioural rhythms may also arise in many other social settings (e.g. in the context of mating interactions²⁹, vigilance behaviour during group foraging). Essentially, the reported diversity suggests that the expectation that individuals within a pair (or group) should optimize their behavioural rhythms relative to the 24-h day may be too simplistic, opening up a wide field to study the evolutionary ecology of plasticity in circadian clocks.

Online Content Methods, along with any additional Extended Data display items and Source Data, are available in the online version of the paper; references unique to these sections appear only in the online paper.



Figure 1 | Map of shorebird breeding sites with data on incubation rhythms. The colour of the dots indicates the genus (data from multiple species per genus may be available), the size of the dots refers to data quality (large: exact breeding

145 site known, small: breeding site estimated, see Methods). For nearby or overlapping locations, the dots are scattered to
 146 increase visibility. Numbers refer to those breeding sites for which incubation rhythms (of one or more species) are
 147 depicted in Fig. 2.




Figure 2 | Actograms illustrating the diversity of shorebird incubation rhythms. **a-b**, Each actogram depicts the bouts of female (yellow; ♀) and male (blue-grey; ♂) incubation at a single nest over a 24-h period, plotted twice, such that each row represents two consecutive days. If present, twilight is indicated by light grey bars (▨) and corresponds to the time when the sun is between 6° and 0° below the horizon, night is indicated by dark grey bars (■) and corresponds to the time when the sun is < 6° below the horizon. Twilight and night are omitted in the centre of the actogram (24:00) to make the incubation rhythm visible. The circled numbers (1 2 3 ...) indicate the breeding site of each pair and correspond to the numbers on the map (Fig. 1). **a**, Between-species diversity. **b**, Within-species diversity. Note that the three rhythms for Western sandpiper and Ringed plover come from the same breeding location.




Figure 3 | Between- and within-species variation in incubation rhythms and their estimated evolution. **a-b**, Box plots are ordered by species (within suborder) from the shortest to the longest median bout length, and depict the genus (colour as in Fig. 1), median (vertical line inside the box), the 25th and 75th percentiles (box), 25th percentiles minus 1.5 times interquartile range and 75th percentile plus 1.5 times interquartile range or minimum/maximum value, whichever is smaller (bars), and the outliers (circles). $N_{\text{median bout length}} = 729$ and $N_{\text{period}} = 584$ nests. **b**, The red vertical line indicates a 24-h period. **c**, Observed and reconstructed incubation bout and period length visualised (by colour) on the phylogenetic tree³⁰ using species' medians (based on population medians) and one of 100 sampled trees (see Methods). The grey circles represent phylogenetically independent contrasts³¹ and hence emphasize the differences at each tree node.




Figure 4 | Relationship between bout and period length. Each dot represents a single nest ($N = 584$ nests), colours depict the genus. In the suborder Scolopaci the median bout length and period length correlate positively ($r_{\text{Spearman}} = 0.56$, $N = 424$ nests); in the suborder Charadrii periods longer than ~ 24 h are absent, and there is no simple relationship between bout and period length ($N = 160$ nests). For species-specific relationships see Extended Data Fig. 2.




Figure 5 | Predictors of variation in incubation rhythms. **a-c**, Relationships between bout length and body size, measured as female wing length (**a**), breeding latitude (**b**) and anti-predation strategy, quantified as escape distance (**c**) for $N = 729$ nest from 91 populations belonging to 32 species. **d**, The relationship between the proportion of nests with a period length that is entrainable by the 24-h light-dark cycle (i.e. period lengths: 3, 6, 12, 24, or 48h) and breeding latitude ($N = 584$ nests from 88 populations belonging to 30 species). **e**, The distribution of period length over latitude. The period was standardized to 24h so that all 24-h harmonics are depicted as 24h (red line) and respective deviations from each harmonic as deviations from 24h (e.g. period of 12.5h is depicted as 25h). **a-e**, Each circle represents the population median; circle size indicates the number of nests. **a-c**, The solid lines depict the model-predicted relationships, the dotted lines the 95% credible intervals based on the joint posterior distribution of 100 separate MCMCglmm runs each with one of the 100 phylogenetic trees with $\sim 1,100$ independent samples per tree. The grey areas depict the predicted relationships for each of 100 runs (i.e. the full range of regression line estimates across 100 models) and illustrate the uncertainty due to the

phylogenetic tree. The predicted relationships stem from a Gaussian phylogenetic mixed-effect models, where the effects of other predictors were kept constant (**a-c**, Extended Data Table 3), or from a binomial phylogenetic mixed-effect model (**d**, Extended Data Table 3).

METHODS

Recording incubation. Incubation data were obtained between 1994 and 2015, for as many shorebird species ($N = 32$) and populations ($N = 91$) as possible, using six methods (for specifications of the equipment see Extended Data Table 4). (1) In 261 nests, a radio frequency identification reader ('RFID') registered presence of tagged parents at the nest. The passive-integrated tag was either embedded in a plastic flag^{32,33}, with which the parents were banded, or glued to the tail feathers³⁴. In 200 nests the RFID was combined with a temperature probe placed between the eggs. The temperature recordings allowed us to identify whether a bird was incubating even in the absence of RFID readings; an abrupt change in temperature demarcated the start or end of incubation³². (2) For 396 nests, light loggers were mounted to the plastic flag or a band that was attached to the bird's leg^{35,36}. The logger recorded maximum light intensity (absolute or relative) for a fixed sampling interval (2-10 min). An abrupt change in light intensity (as opposed to a gradual change caused, e.g. by civil twilight) followed by a period of low or high light intensity demarcated the start or end of the incubation period (Extended Data Fig. 6). (3) For nine nests a GPS tag, mounted on the back of the bird, recorded the position of the bird³⁷. The precision of the position depends on cloud cover and sampling interval³⁷. Hence, to account for the imprecision in GPS positions, we assumed incubation whenever the bird was within 25 m of the nest (Extended Data Fig. 6b). (4) At three nests automated receivers recorded signal strength of a radio-tag attached to the rump of a bird; whenever a bird incubated, the strength of the signal remained constant²⁶ (Supplementary Actograms p. 257-9¹⁸). (5) At 53 nests video cameras and (6) for 8 nests continuous observations were used to identify the incubating parents; parent identification was based on plumage, colour rings or radio-tag. In one of the populations, three different methods were used, in seven populations representing seven species two methods. In one nest, two methods were used simultaneously (Extended Data Fig. 6b).

Extraction of incubation bouts. An incubation bout was defined as the total time allocated to a single parent (i.e. the time between the arrival of a parent at and its departure from the nest followed by incubation of its partner). Bout lengths were only extracted if at least 24h of continuous recording was available for a nest; in such cases, all bout lengths were extracted. For each nest, we transformed the incubation records to local time as $(\text{UTC time} + \frac{\text{nest's longitude}}{15})$. Incubation bouts from RFIDs, videos and continuous observations were mostly extracted by an R-script and the results verified by visualizing the extracted and the raw data^{18,32,38,39}; otherwise, MB extracted the bouts manually from plots of raw data^{40,41} (plots of raw data and extracted bouts for all nests are in the Supplementary Actograms¹⁸; the actograms were generated by 'ggplot' and 'xyplot' functions from the 'ggplot2' and 'lattice' R package⁴²⁻⁴⁴). Whenever the start or end of a bout was unclear, we classified these bouts as uncertain (see next paragraph for treatment of unsure bouts). In case of light logger data, the light recordings before and after the breeding period, when the birds were definitely not incubating, helped to distinguish incubation from non-incubation. Whenever an individual tagged with a light logger nested in an environment where the sun was more than 6° below the horizon for part of a day (i.e. night), we assumed an incubation bout when the individual started incubating before the night started and ended incubating after the night ended. When different individuals incubated at the beginning vs. at the end of the night, we either did not quantify these bouts or we indicated the possible time of exchange (based on trend in previous exchanges), but classified these bouts as uncertain (see Supplementary Actograms¹⁸). In total, we extracted 34,225 incubation bouts.

The proportion of uncertain bouts within nests had a distribution skewed towards zero (median = 0%, range: 0-100%, $N = 729$ nests), and so did the median proportion of uncertain bouts within populations (median = 2%, range: 0-74%, $N = 91$ populations). Excluding the uncertain bouts did not change our estimates of median bout length (Pearson's correlation coefficient for median bout length based on all bouts and without uncertain bouts: $r = 0.96$, $N = 335$ nests with both certain and uncertain bouts). Hence, in further analyses all bouts were used to estimate median bout length.

Note that in some species sexes consistently differ in bout length (Figure 2a, e.g. Northern lapwing). As these differences are small compared to the between-species differences and because in 27 nests (of 8 species) the sex of the parents was unknown, we here use median bout length independent of sex.

Extraction of period length. The method used for extracting the period length of incubation rhythm for each nest is described in the Extended Data Fig. 1.

Extraction of entrainable periods. We classified 24-h periods and periods with 24-h harmonics (i.e. 3, 6, 12, 48h) as strictly entrainable by 24-h light fluctuations ($N = 142$ nests out of 584). Including also nearest adjacent periods (± 0.25 h) increased the number of nests with entrainable periods ($N = 277$), but results of statistical analyses remained quantitatively similar. We consider periods and harmonics of 12.42h (i.e. 3.1, 6.21, 12.42, 24.84h) as strictly entrainable by tide. However, because the periods in our data were extracted in 0.25-h intervals (Extended Data Fig. 1), we classified periods of 3, 6.25, 12.5, 24.75h (i.e. those closest to the strict tide harmonics) as entrainable by tide ($N = 32$ nests out of 584). Including also the second nearest periods (i.e. 3.25, 6, 12.25, 25) increased the number of nests entrainable by tide to $N = 55$.

Population or species life-history traits. For 643 nests, the exact breeding location was known (nests or individuals were monitored at the breeding ground). For the remaining 86 nests (from 27 populations representing 8 species, where individuals were tagged with light loggers on the wintering ground), the breeding location was roughly estimated from the recorded 24-h variation in daylight, estimated migration tracks, and the species' known breeding range⁴⁵⁻⁵². One exact breeding location was in the Southern Hemisphere, so we used absolute latitude in analyses. Analyses without populations with estimated breeding-location or without the Southern Hemisphere population generated quantitatively similar estimates as the analyses on full data.

For each population, body size was defined as mean female wing length⁵³, either for individuals measured at the breeding area or at the wintering area. In case no individuals were measured, we used the mean value from the literature (see open access data for specific values and references⁵⁴).

Anti-predation strategy was assessed by estimating escape distance of the incubating bird when a human approached the nest, because species that are cryptic typically stay on the nest much longer than non-cryptic species, sometimes until nearly stepped upon^{49,55}. Escape distance was obtained for all species. Forty-

four authors of this paper estimated the distance (in m) for one or more species based on their own data or experience. For 10 species, we also obtained estimates from the literature⁴⁹. We then used the median ‘estimated escape distance’ for each species. In addition, for 13 species we obtained ‘true escape distance’. Here, the researcher approached a nest (of known position) and either estimated his distance to the nest or marked his position with GPS when the incubating individual left the nest. For each GPS position, we calculated the Euclidian distance from the nest. In this way we obtained multiple observations per nest and species, and we used the median value per species (weighted by the number of estimates per nest) as the ‘true escape distance’. The species’ median ‘estimated escape distance’ was a good predictor of the ‘true escape distance’ (Pearson’s correlation coefficient: $r = 0.89$, $N = 13$ species). For analysis, we defined the escape distance of a species as the median of all available estimates.

For each species, we determined whether it predominantly uses a tidal environment outside its breeding ground, i.e. has tidal vs. non-tidal life history (based on^{49,51,52}). For each population with exact breeding location, we scored whether tidal foraging habitats were used by breeding birds for foraging (for three populations this information was unknown)⁵⁴. For all populations with estimated breeding location we assumed, based on the estimated location and known behaviour at the breeding grounds, no use of tidal habitat.

Statistical analyses. Unless specified otherwise, all analyses were performed on the nest level using median bout length and extracted period length.

We used phylogenetically informed comparative analyses to assess how evolutionary history constrains the incubation rhythms (estimated by Pagel’s λ coefficient of phylogenetic signal^{56,57}) and to control for potential non-independence among species due to common ancestry. This method explicitly models how the covariance between species declines as they become more distantly related^{56,58,59}. We used the Hackett⁶⁰ backbone phylogenetic trees available at <http://birdtree.org>⁶¹, which included all but one species (*Charadrius nivosus*) from our dataset. Following a subsequent taxonomic split⁶², we added *Charadrius nivosus* to these trees as a sister taxon of *Charadrius alexandrinus*. Phylogenetic uncertainty was accounted for by fitting each model with 100 phylogenetic trees randomly sampled from 10,000 phylogenies at <http://birdtree.org>⁶¹.

The analyses were performed with Bayesian phylogenetic mixed-effect models (Fig. 5 and Extended Data Table 1 and 3) and the models were run with the ‘MCMCglmm’ function from the R package ‘MCMCglmm’⁶³. In all models, we also accounted for multiple sampling within species and breeding site (included as random effects). In models with a Gaussian response variable, an inverse-gamma prior with shape and scale equal to 0.001 was used for the residual variance (i.e. variance set to one and the degree of belief parameter to 0.002). In models with binary response variables, the residual variance was fixed to one. For all other variance components the parameter-expanded priors were used to give scaled F-distributions with numerator and denominator degrees of freedom set to one and a scale parameter of 1,000. Model outcomes were insensitive to prior parameterization. The MCMC chains ran for 2,753,000 iterations with a burn-in of 3,000 and a thinning interval of 2,500. Each model generated ~1,100 independent samples of model parameters (Extended Data Table 1 and 3). Independence of samples in the Markov chain was assessed by tests for autocorrelation between samples and by using graphic diagnostics.

First, we used MCMCglmm to estimate Pagel’s λ (phylogenetic signal) for bout and period length (Gaussian), and to show that our estimates of these two incubation variables were independent of how often the incubation behaviour was sampled (‘sampling’ in min, ln-transformed; Extended Data Table 1). Hence, in subsequent models, sampling was not included.

Then, we used MCMCglmm to model variation in bout length and period length. Bout length was modelled as a continuous response variable and latitude (°, absolute), female wing length (mm, ln-transformed) and approach distance (m, ln-transformed) as continuous predictors. Predictors had low collinearity (at nest, population and species level; all Pearson or Spearman correlation coefficients $|r| < 0.28$). To test for potential entrainment to 24-h, period length was modelled as a binary response variable (1 = rhythms with period of 3, 6, 12, 24, or 48 h; 0 = rhythms with other periods) and latitude as a continuous predictor. To test how circadian period varies with latitude or life history, period was transformed to deviations from 24-h and 24-h harmonics and scaled by the time span between the closest harmonic and the closest midpoint between two harmonics. For example, a 42h period deviates by -6h from 48h (the closest 24-h harmonic) and hence -6h was divided by 12h (the time between 36h – the midpoint of two harmonics - and 48h -the closest harmonic). This way the

deviations spanned from -1 to 1 with 0 representing 24-h and its harmonics. The absolute deviations were then modelled as a continuous response variable and latitude as continuous predictor. The deviations were also modelled as a continuous response and species life history (tidal or not) as categorical predictor.

In all models the continuous predictor variables were centred and standardized to a mean of zero and a standard deviation of one.

We report model estimates for fixed and random effects, as well as for Pagel's λ , by the modes and the uncertainty of the estimates by the highest posterior density intervals (referred to as 95% CI) from the joint posterior distributions of all samples from the 100 separate runs, each with one of the 100 separate phylogenetic trees from <http://birdtree.org>.

To help interpret the investigated relationships we assessed whether incubation rhythms evolved within diverged groups of species by plotting the evolutionary tree of the incubation rhythm variables (Fig. 3b), as well as of the predictors (Extended Data Fig. 5).

The source of phylogenetic constraint in bout and period length was investigated by estimating the proportion of phenotypic variance explained by suborder, genus and species (Extended Data Table 2). The respective mixed models were also specified with 'MCMCglmm'⁶³ using the same specifications as in the phylogenetic models. Because suborder contained only two levels, we first fitted an intercept mixed model with genus, species, and breeding site as random factors, and used it to estimate the overall phenotypic variance. We then entered suborder as a fixed factor and estimated the variance explained by suborder as the difference between the total variance from the first and the second model. To evaluate the proportion of the variance explained by species, genus and breeding site, we used the estimates from the model that included suborder.

R version 3.1.1⁶⁴ was used for all statistical analyses.

Code availability. All statistical analyses are replicable with the open access data and r-code available from <https://osf.io/wxufm/> (not public yet, but available for peer review: https://osf.io/wxufm/?view_only=d0172c5f0935462f9471d6fdc4e40d86).

REFERENCES

- 1 Dunlap, J. C., Loros, J. J. & DeCoursey, P. J. *Chronobiology: Biological Timekeeping* (Sinauer Associates, 2004).
- 2 Kronfeld-Schor, N., Bloch, G. & Schwartz, W. J. Animal clocks: when science meets nature. *Proc R Soc B* **280** (2013).
- 3 Bloch, G., Herzog, E. D., Levine, J. D. & Schwartz, W. J. Socially synchronized circadian oscillators. *Proc R Soc B* **280** (2013).
- 4 Davidson, A. J. & Menaker, M. Birds of a feather clock together—sometimes: social synchronization of circadian rhythms. *Curr Opin Neurobiol* **13**, 765-769 (2003).
- 5 Castillo-Ruiz, A., Paul, M. J. & Schwartz, W. J. In search of a temporal niche: social interactions. *Prog Brain Res* **199**, 267-280 (2012).
- 6 Emlen, S. T. & Oring, L. W. Ecology, sexual selection, and the evolution of mating systems. *Science* **197**, 215-223 (1977).
- 7 Young, M. W. & Kay, S. A. Time zones: a comparative genetics of circadian clocks. *Nat Rev Genet* **2**, 702-715 (2001).
- 8 Helm, B. & Visser, M. E. Heritable circadian period length in a wild bird population. *Proc R Soc B* (2010).
- 9 Koskenvuo, M., Hublin, C., Partinen, M., Heikkilä, K. & Kaprio, J. Heritability of diurnal type: a nationwide study of 8753 adult twin pairs. *J Sleep Res* **16**, 156-162 (2007).
- 10 Mrosovsky, N. Locomotor activity and non-photoc influences on circadian clocks. *Biol Rev* **71**, 343-372 (1996).
- 11 Mistlberger, R. E. & Skene, D. J. Social influences on mammalian circadian rhythms: animal and human studies. *Biol Rev* **79**, 533-556 (2004).
- 12 Regal, P. J. & Connolly, M. S. Social influences on biological rhythms. *Behaviour* **72**, 171-198 (1980).
- 13 Schoener, T. W. Resource partitioning in ecological communities. *Science* **185**, 27-39 (1974).
- 14 Paul, M. J., Indic, P. & Schwartz, W. J. Social synchronization of circadian rhythmicity in female mice depends on the number of cohabiting animals. *Biol Lett* **11**, 20150204 (2015).

360 15 Deeming, D. C. *Avian Incubation: Behaviour, Environment and Evolution* (Oxford University Press, 2002).

361 16 Szekely, T. & Reynolds, J. D. Evolutionary transitions in parental care in shorebirds. *Proc R Soc B* **262**, 57-64

362 (1995).

363 17 del Hoyo, J., Elliott, A. & Sargatal, J. *Handbook of the Birds of the World. Vol. 3. Hoatzing to Auks*. (Lynx

364 Edicions, 1996).

365 18 Bulla, M. *et al.* Supplementary Actograms from Defying the 24-h day: Unexpected diversity in socially

366 synchronized rhythms of shorebirds. Version 1. *Open Science Framework*,

367 https://osf.io/9dtpa/?view_only=3c5fbf6d176e41afb83c4306604e9e10 (2016).

368 19 Williams, J. B. in *Avian Energetics and Nutritional Ecology* (ed C. Carey) Ch. 5, 375-416 (Chapman & Hall,

369 1996).

370 20 Bulla, M., Cresswell, W., Rutten, A. L., Valcu, M. & Kempenaers, B. Biparental incubation-scheduling: no

371 experimental evidence for major energetic constraints. *Behav Ecol* **26**, 30-37 (2015).

372 21 Martin, T. E., Scott, J. & Menge, C. Nest predation increases with parental activity: separating nest site and

373 parental activity effects. *Proc R Soc B* **267**, 2287-2293 (2000).

374 22 Smith, P. A., Tulp, I., Schekkerman, H., Gilchrist, H. G. & Forbes, M. R. Shorebird incubation behaviour and

375 its influence on the risk of nest predation. *Anim Behav* **84**, 835-842 (2012).

376 23 McKinnon, L. *et al.* Lower predation risk for migratory birds at high latitudes. *Science* **327**, 326-327 (2010).

377 24 Hut, R. A., Paolucci, S., Dor, R., Kyriacou, C. P. & Daan, S. Latitudinal clines: an evolutionary view on

378 biological rhythms. *Proc R Soc B* **280**, 20130433 (2013).

379 25 van Oort, B. E. *et al.* Circadian organization in reindeer. *Nature* **438**, 1095-1096 (2005).

380 26 Steiger, S. S. *et al.* When the sun never sets: diverse activity rhythms under continuous daylight in free-

381 living arctic-breeding birds. *Proc R Soc B* **280** (2013).

382 27 Foster, R. G. & Wulff, K. The rhythm of rest and excess. *Nat Rev Neurosci* **6**, 407-414 (2005).

383 28 Silver, R. & Bittman, E. L. Reproductive Mechanisms: Interaction of Circadian and Interval Timing. *Ann N Y*

384 *Acad Sci* **423**, 488-514 (1984).

385 29 Lesku, J. A. *et al.* Adaptive sleep loss in polygynous pectoral sandpipers. *Science* **337**, 1654-1658 (2012).

386 30 Revell, L. J. & Freckleton, R. Two new graphical methods for mapping trait evolution on phylogenies.
387 *Methods. Ecol. Evol.* **4**, 754-759, <http://dx.doi.org/10.1111/2041-210x.12066> (2013).

388 31 Felsenstein, J. Phylogenies and the comparative method. *Am Nat* **125**, 1-15 (1985).

389 32 Bulla, M., Valcu, M., Rutten, A. L. & Kempenaers, B. Biparental incubation patterns in a high-Arctic breeding
390 shorebird: how do pairs divide their duties? *Behav Ecol* **25**, 152-164 (2014).

391 33 Reneerkens, J., Grond, K., Schekkerman, H., Tulp, I. & Piersma, T. Do uniparental Sanderlings *Calidris alba*
392 increase egg heat input to compensate for low nest attentiveness? *PLoS ONE* **6**, e16834,
393 <http://dx.doi.org/10.1371/journal.pone.0016834> (2011).

394 34 Kosztolányi, A. & Székely, T. Using a transponder system to monitor incubation routines of Snowy Plovers. *J*
395 *Field Ornithol* **73**, 199-205 (2002).

396 35 Conklin, J. R. & Battley, P. F. Attachment of geolocators to bar-tailed godwits: a tibia-mounted method with
397 no survival effects or loss of units. *Wader Study Group Bull* **117**, 56-58 (2010).

398 36 Burger, J., Niles, L. J., Porter, R. R. & Dey, A. D. Using geocator data to reveal incubation periods and
399 breeding biology in Red Knots *Calidris canutus rufa*. *Wader Study Group Bull* **119**, 26-36 (2012).

400 37 Bouten, W., Baaij, E. W., Shamoun-Baranes, J. & Camphuysen, K. C. J. A flexible GPS tracking system for
401 studying bird behaviour at multiple scales. *J Ornithol* **154**, 571-580 (2012).

402 38 Bulla, M. R-SCRIPT and EXAMPLE DATA to extract incubation from temperature measurements. Version 1.
403 *figshare*, <https://dx.doi.org/10.6084/m9.figshare.1037545.v1> (2014).

404 39 Bulla, M. R-SCRIPT and EXAMPLE DATA to extract incubation bouts from continuous RFID and video data.
405 Version 1. *figshare*, <http://dx.doi.org/10.6084/m9.figshare.1533278.v1> (2015).

406 40 Bulla, M. Example of how to manually extract incubation bouts from interactive plots of raw data - R-CODE
407 and DATA. Version 1. *figshare*, <https://dx.doi.org/10.6084/m9.figshare.2066784.v1> (2016).

408 41 Bulla, M. Procedure for manual extraction of incubation bouts from plots of raw data.pdf. Version 1.
409 *figshare*, <https://dx.doi.org/10.6084/m9.figshare.2066709.v1> (2016).

410 42 Wickham, H. *ggplot2: Elegant Graphics for Data Analysis* (Springer Publishing Company, Incorporated,
411 2009).

412 43 Sarkar, D. & Andrews, F. latticeExtra: Extra Graphical Utilities Based on Lattice. R package version 0.6-24.
413 <http://CRAN.R-project.org/package=latticeExtra> (2012).

414 44 Sarkar, D. *Lattice: Multivariate Data Visualization with R*. (Springer, 2008).

415 45 Lisovski, S. Geolocator-ArcticWader-BreedingSiteEstimation. Version 2015-08-05. *GitHub repository*,
416 <https://github.com/slisovski/Geolocator-ArcticWader-BreedingSiteEstimation> (2015).

417 46 Lisovski, S. & Hahn, S. GeoLight – processing and analysing light-based geolocator data in R. *Methods. Ecol.*
418 *Evol.* **3**, 1055-1059, <https://dx.doi.org/10.1111/j.2041-210X.2012.00248.x> (2012).

419 47 Lisovski, S. *et al.* Geolocation by light: accuracy and precision affected by environmental factors. *Methods.*
420 *Ecol. Evol.*, 1-10, <https://dx.doi.org/10.1111/j.2041-210X.2012.00185.x> (2012).

421 48 Conklin, J. R., Battley, P. F., Potter, M. A. & Fox, J. W. Breeding latitude drives individual schedules in a
422 trans-hemispheric migrant bird. *Nat Commun* **1**, 67 (2010).

423 49 Poole, A. *The Birds of North America Online* (Cornell Laboratory of Ornithology, 2005).

424 50 Lappo, E., Tomkovich, P. & Syroechkovskiy, E. *Atlas of Breeding Waders in the Russian Arctic* (UF Ofsetnaya
425 Pecha, 2012).

426 51 Chandler, R. J. *Shorebirds of the Northern Hemisphere* (Christopher Helm, 2009).

427 52 Brazil, M. *Birds of East Asia: Eastern China, Taiwan, Korea, Japan, and Eastern Russia* (Christohper Helm,
428 2009).

429 53 Dale, J. *et al.* Sexual selection explains Rensch's rule of allometry for sexual size dimorphism. *Proc R Soc B*
430 **274**, 2971-2979 (2007).

431 54 Bulla, M. *et al.* Supplementary Data 3 - Study sites: location, population wing length, monitoring method,
432 tide. Version 11. *figshare*, <https://dx.doi.org/10.6084/m9.figshare.1536260.v11> (2016).

433 55 Cramp, S. *Handbook of the Birds of Europe, the Middle East, and North Africa: The Birds of the Western*
434 *Palaearctic Volume III: Waders to Gulls* (Oxford University Press, 1990).

435 56 Freckleton, R. P., Harvey, P. H. & Pagel, M. Phylogenetic analysis and comparative data: a test and review of
436 evidence. *Am Nat* **160**, 712-726 (2002).

437 57 Pagel, M. Inferring evolutionary processes from phylogenies. *Zool Scr* **26**, 331-348 (1997).

438 58 Martins, E. P. & Hansen, T. F. Phylogenies and the comparative method: a general approach to
439 incorporating phylogenetic information into the analysis of interspecific data. *Am Nat*, 646-667 (1997).

440 59 Pagel, M. Inferring the historical patterns of biological evolution. *Nature* **401**, 877-884 (1999).

441 60 Hackett, S. J. *et al.* A phylogenomic study of birds reveals their evolutionary history. *Science* **320**, 1763-1768
442 (2008).

443 61 Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K. & Mooers, A. O. The global diversity of birds in space and
444 time. *Nature* **491**, 444-448 (2012).

445 62 Küpper, C. *et al.* Kentish versus Snowy Plover: phenotypic and genetic analyses of *Charadrius alexandrinus*
446 reveal divergence of Eurasian and American subspecies. *Auk* **126**, 839-852 (2009).

447 63 Hadfield, J. D. MCMC methods for multi-response generalized linear mixed models: the MCMCglmm R
448 package. *J Stat Softw* **33**, 1-22 (2010).

449 64 R-Core-Team. R: A Language and Environment for Statistical Computing. Version 3.1.1. *R Foundation for*
450 *Statistical Computing*, <http://www.R-project.org/> (2014).

451 65 Anderson, D. R. *Model Based Inference in the Life Sciences: A Primer on Evidence* (Springer, 2008).

452 66 Hijmans, R. J. raster: Geographic data analysis and modeling. R package version 2.3-24. [http://CRAN.R-](http://CRAN.R-project.org/package=raster)
453 [project.org/package=raster](http://CRAN.R-project.org/package=raster) (2015).

454 67 Bivand, R. & Lewin-Koh, N. maptools: Tools for reading and handling spatial objects. R package version 0.8-
455 30. <http://CRAN.R-project.org/package=maptools> (2014).

456 68 Revell, L. J. in *Modern Phylogenetic Comparative Methods and Their Application in Evolutionary Biology* (ed
457 L. Z. Garamszegi) Ch. 4, 77-103 (Springer, 2014).

458 69 Orme, D. *et al.* caper: Comparative Analyses of Phylogenetics and Evolution in R. R package version 0.5.2.
459 <http://CRAN.R-project.org/package=caper> (2013).

460 70 Johnson, O. W. *et al.* Tracking Pacific Golden-Plovers *Pluvialis fulva*: transoceanic migrations between non-
461 breeding grounds in Kwajalein, Japan and Hawaii and breeding grounds in Alaska and Chukotka. *Wader*
462 *Study* **122**, 13-20 (2015).

- 463 71 Kosztolanyi, A., Cuthill, I. C. & Székely, T. Negotiation between parents over care: reversible compensation
 464 during incubation. *Behav Ecol* **20**, 446-452 (2009).
- 465 72 St Clair, J. J. H., Herrmann, P., Woods, R. W. & Székely, T. Female-biased incubation and strong diel sex-roles
 466 in the Two-banded Plover *Charadrius falklandicus*. *J Ornithol*, 1-6 (2010).
- 467 73 Spiegel, C. S., Haig, S. M., Goldstein, M. I. & Huso, M. Factors affecting incubation patterns and sex roles of
 468 Black Oystercatchers in Alaska. *Condor* **114**, 123-134 (2012).
- 469 74 Praus, L. & Weidinger, K. Predators and nest success of Sky Larks *Alauda arvensis* in large arable fields in the
 470 Czech Republic. *Bird Study* **57**, 525-530 (2010).

471 **Supplementary Information** is linked to the online version of the paper at www.nature.com/nature.


472 **Acknowledgements** We thank all that made the data collection possible. We are grateful to W. Schwartz, E. Schlicht, W.
 473 Forstmeier, M. Baldwin, D. Starr-Glass, and B. Bulla for comments on the manuscript and to F. Korner-Nievergelt, J. D.
 474 Hadfield, L. Z. Garamszegi, S. Nakagawa, T. Roth, N. Dochtermann, Y. Araya, E. Miller and H. Schielzeth for advice on data
 475 analysis. Data collection was supported by various institutions and people listed in the Supplementary Data 1. The study
 476 was supported by the Max Planck Society (to B.K.). M.B. is a PhD student in the International Max Planck Research School
 477 for Organismal Biology.

478 **Author Contributions** M.B. and B.K. conceived the study. All authors except for B.H. collected the primary data. MB
 479 managed the data. MB and M.V. developed the methods to extract incubation. M.B. extracted bout lengths and with help
 480 from A.R. and M.V. created actograms. M.B. with help from M.V. analysed the data. M.B. and B.K. wrote the paper with
 481 input from the other authors. Except for the first, second and last author, the authors are listed alphabetically by their first
 482 name.

483 **Author Information** All information, primary and extracted data, computer code and software necessary to replicate our
 484 results, as well as the Supplementary Actograms are open access and archived at Open Science Framework
 485 <https://osf.io/wxufm/> (not public yet, but available for peer review: https://osf.io/wxufm/?view_only=d0172c5f0935462f9471d6fdc4e40d86). Reprints and permissions information is available at www.nature.com/reprints.
 487 The authors declare no competing financial interests. Readers are welcome to comment on the online version of the paper.
 488 Correspondence and requests for materials should be addressed to M.B. (bulla.mar@gmail.com) and B.K.

489 (b.kempenaers@orn.mpg.de).

490 EXTENDED DATA



491

492

493

494

495

496

497

498

499

500

501

502

503


504


505

506


Extended Data Figure 1 | Illustration of the method used for extracting period length of incubation rhythm. **a-c**, Each column represents an example for a specific nest with long, intermediate and short incubation bouts. **a**, From the extracted bout lengths we created a time series that indicated for each nest, and every 10 min, whether a specific parent (female, if sex was known) incubated or not. Exchange gaps (no parent on the nest) had to be < 6 h to be included (for treatment of exchange gaps > 6 h see **d**, **e**). **b**, We then estimated the autocorrelation for each 10 min time-lag up to 4 days (R 'acf' function⁶⁴). Positive values indicate a high probability that the female was incubating, negative values indicate that it was more likely that the male was incubating. We used only nests that had enough data to estimate the autocorrelation pattern ($N = 584$ nests from 88 populations of 30 species). The visualized autocorrelation time series never resembled white or random noise indicative of an arrhythmic incubation pattern. To determine the period (i.e. cycle of high and low probability for a parent to incubate) that dominated the incubation rhythm, we fitted to the autocorrelation estimates a series of periodic logistic regressions. In each regression, the time lag (in hours) transformed to radians was represented by a sine and cosine function $f(t) = a_0 + b \cdot \left(\cos \frac{2 \cdot \pi \cdot t}{T} \right) + c \cdot \left(\sin \frac{2 \cdot \pi \cdot t}{T} \right) + e$, where $f(t)$ is the autocorrelation at time-lag t , a_0 is the intercept, b is the slope for sine and c the slope for cosine, T represents the length of the fitted period (in hours), and e is an error term. We allowed the period length to vary from 0.5 h to 48 h (in 15 min intervals, giving 191 regressions). **c**, By comparing the Akaike's Information Criterion⁶⁵ (AIC) of all regressions, we estimated, for each nest, the length of the dominant period in the actual incubation data (best fit). Regressions with ΔAIC ($AIC_{\text{model}} - AIC_{\text{min}}$) close to 0 are considered as having strong empirical support, while models with ΔAIC values ranging from 4-7 have less

support⁶⁵. In 73% of all nests, we determined a single best model with $\Delta AIC \leq 3$ (**c**, middle ΔAIC graph), in 20% of nests two best models emerged and in 6% of nests 3 or 4 models had $\Delta AIC \leq 3$ (**c**, left and right ΔAIC graphs). However, in all but three nests, the models with the second, third, etc. best ΔAIC were those with period lengths closest to the period length of the best model (**c**, left and right ΔAIC graphs). This suggests that multiple periodicities are uncommon. **d-e**, The extraction of the period length (described in **a-c**) requires continuous datasets, but some nests had long (>6 h) gaps between two consecutive incubation bouts, for example because of equipment failure or because of unusual parental behaviour. In such cases, we excluded the data from the end of the last bout until the same time the following day, if data were then available again (**d**), or we excluded the entire day (**e**).






Extended Data Figure 3 | Ecological correlates of latitude. **a**, Variation in minimum temperature across the globe represented by mean minimum June temperature for the Northern Hemisphere and mean minimum December temperature for the Southern Hemisphere. **b**, Correlation between absolute latitude and the mean minimum temperature based on the month represented by mid-day of incubation data for each nest ($N = 729$). For maximum temperature the correlation was similar ($r = -0.91$, $N = 729$ nests). **c**, Daily variation in sun elevation (i.e. in light conditions) represented as the difference between the noon and midnight sun-elevation for the summer solstice in the Northern Hemisphere and the winter solstice in the Southern Hemisphere. **d**, Correlation between absolute latitude and daily variation in sun elevation for mid-day of incubation data for each nest ($N = 729$ nests). The points are jittered, as else they form a straight line. **a, c**, Red points indicate the breeding sites ($N = 91$). **a-b**, The minimum and maximum monthly temperature data were obtained from www.worldclim.org using the 'raster' R-package⁶⁶. **c-d**, Sun-elevation was obtained by the 'solarpos' function from the 'maptools' R-package⁶⁷.




528
529
530
531
532
533

Extended Data Figure 4 | Between-species variation in parental crypsis during incubation. **a-b**, Shorebirds vary in how visible they are on the nest while incubating. The nearly invisible Great knot (*Calidris tenuirostris*; **a**; central and facing right) sits tight on the nest when approached by a human until nearly stepped upon. In contrast, the conspicuous Eurasian oystercatcher (*Haematopus ostralegus*; **b**) is visible on the nest from afar and when approached by a human leaves the nest about 100 m in advance (Credits: **a**, M. Šálek; **b**, Jan van de Kam).



Extended Data Figure 5 | Phylogenetic relationships for predictors. a, Body size, estimated as female wing length. **b**, Latitude (absolute), **c**, Escape distance. **a-c**, We visualised the evolution of these traits^{30,68} using species' medians (**a-b**; based on population medians), species' estimates of escape distance (**c**) and one of the 100 sampled trees (see Methods).



Extended Data Figure 6 | Illustration and verification of the method used to extract incubation bouts from light logger data. a, An example of a nest with a light intensity signal from both parents (yellow line — and blue-grey line —). The incubation bouts for a given parent reflect periods dominated by lower light values compared to those of the partner. Note the sharp drop in the light levels at the beginning of each incubation bout and the sharp increase in the light levels at the end. Change-overs between partners occur when the light signal lines cross. Such pronounced changes in light intensity detected by the logger were used to assign incubation even when only a single parent was tagged. Note that after the chicks hatch and leave the nest (July 9, vertical bar), the light intensity signal from both parents remains similar. **b,** An example of a nest where one incubating parent was simultaneously equipped with a light logger and with a GPS tag. The yellow line (—) indicates light levels, red dots (•) indicate the distance of the bird to the nest. As expected, low light levels co-occur with close proximity to the nest, and hence reflect periods of incubation. Although light levels decrease during twilight (light grey horizontal bar; ■), the recordings were still sensitive enough to reflect periods of incubation, i.e. the light signal matches the distance (e.g. May-25: female incubated during dawn, but was off the nest during dusk). **a-b,** Rectangles in the background indicate extracted female (light yellow polygon, ■) and male (light blue-grey polygon, ■) incubation bouts.

Extended Data Table 1 | Effects of phylogeny and sampling on bout length and period length.

Response	Effect type	Effect	Posterior mode	95% CI		N (range)
				Lower	Upper	
Median bout [h]	Fixed	Intercept	7.2	1.04	12	1100 (924-2079)
		Sampling	0.16	-0.2	0.61	1100 (809-1644)
	Random (variance)	Phylogeny	25.33	4.6	59.6	1100 (753-1383)
		Species	0.01	0	12.1	1100 (779-1636)
		Breeding site	2.13	0.96	4.28	1100 (808-2242)
		Residual	5.04	4.51	5.61	1100 (838-1444)
	Pagel's λ		1	0.5	1	1100 (814-1316)
Period [h]	Fixed	Intercept	21.94	12.8	30.67	1100 (765-1392)
		Sampling	0.13	-0.41	0.65	1100 (741-1468)
	Random (variance)	Phylogeny	66.22	14.3	153	1100 (729-1638)
		Species	0.06	0	29.36	1100 (729-1435)
		Breeding site	0.01	0	0.88	1100 (814-1378)
		Residual	14.87	13.3	16.84	1100 (884-1460)
	Pagel's λ		1	0.54	1	1100 (740-1523)

The posterior estimates (modes) of the effect sizes with the highest posterior density intervals (95% CI) and the median and range of the effective sample sizes (N (range)) come from the joint posterior distribution of 100 separate runs each with one of 100 separate phylogenetic trees from <http://birdtree.org>. $N_{\text{bout}} = 729$ nests from 91 populations belonging to 32 species. $N_{\text{period}} = 584$ nests from 88 populations belonging to 30 species. Sampling (how often the incubation behaviour was sampled) was ln-transformed and then mean-centred and scaled (divided by SD). For procedures and specifications related to phylogenetic Bayesian mixed models see Methods. Estimating Pagel's λ on the species level ($N_{\text{bout}} = 32$ species, $N_{\text{period}} = 30$ species) with phylogenetic generalized least-squares using the function 'pgls' from the R-package 'caper'⁶⁹ gave similar results (median (range) $\lambda_{\text{bout}} = 0.73$ (0.63-1) and $\lambda_{\text{period}} = 0.95$ (0.64-1), based on 100 estimates each for one of the 100 trees).

559 **Extended Data Table 2 | Source of phylogenetic signal**

Suborder included	Response	Effect type	Fixed effects	Posterior mode	95% CI		N
					Lower	Upper	
No	Bout [h]	Fixed	Intercept	7.69	4.11	10.76	1100
			Genus	9.54	0.02	46.6	1100
		Random (variance)	Species	11	4.84	25.77	1100
			Breeding site	2.25	1.06	4.5	1100
			Residual	5.03	4.55	5.61	1100
Yes	Bout [h]	Fixed	Intercept (Charadrii)	4.48	1.05	7.66	1100
			Suborder (Scolopaci)	6.07	1.52	10.44	1100
		Random (variance)	Genus	0.09	0	21.35	1100
			Species	11.29	5.53	23.99	1275
			Breeding site	2.31	0.98	4.28	1100
			Residual	5.04	4.52	5.58	1100
No	Period [h]	Fixed	Intercept	23.46	18.22	27.39	1100
			Genus	19.05	0	97.79	1100
		Random (variance)	Species	41.02	17.04	72.66	1100
			Breeding site	0.01	0	0.9	1100
			Residual	14.94	13.35	16.76	1100
Yes	Period [h]	Fixed	Intercept (Charadrii)	18.61	13.12	23.98	940
			Suborder (Scolopaci)	8.92	0.6	15.4	990
		Random (variance)	Genus	0.33	0	59.09	769
			Species	34.41	16.91	68.14	1142
			Breeding site	0.01	0	0.84	1100
			Residual	15.01	13.34	16.86	891

560 The posterior estimates (modes) of the effect sizes with the highest posterior density intervals (95% CI) and the effective sample sizes (N) come from a
561 posterior distribution of 1,100 simulated values generated by ‘MCMCglmm’ in R⁶³. N_{bout} = 729 nests from 91 populations belonging to 32 species. N_{period} =
562 584 nest from 88 populations belonging to 30 species.

Extended Data Table 3 | Effect of latitude, body size, escape distance and life history on biparental incubation rhythms in shorebirds

Response	Effect type	Fixed effects	Posterior mode	95% CI		N (range)
				Lower	Upper	
Bout [h]	Fixed	Intercept	7.45	2.65	12	1100 (804-1496)
		Wing length	-0.78	-2.5	1.05	1100 (839-1638)
		Latitude	1.72	0.63	2.65	1100 (850-1642)
		Escape distance	-1.68	-3.3	-0.25	1100 (634-2046)
	Random (variance)	Phylogeny	0.19	0	45	1100 (803-1875)
		Species	0.07	0	14.4	1100 (695-1580)
		Breeding site	1.4	0.59	3.02	1100 (833-1480)
		Residual	5.02	4.53	5.64	1100 (516-1916)
	Page's λ		0.72	0.13	1	1100 (731-1407)
Light entrainable rhythm [1,0] on binomial scale	Fixed	Intercept	-1.62	-3.19	-0.13	1100 (731-1633)
		Latitude	-0.56	-1.15	-0.07	1100 (765-1575)
	Random (variance)	Phylogeny	0.05	0	5.54	1100 (883-1371)
		Species	0.02	0	2.68	1100 (965-2246)
		Breeding site	0	0	0.63	1100 (605-1304)
	Page's λ		0.74	0.02	1	1100 (932-1498)
Absolute deviations from 24-h	Fixed	Intercept	0.17	-0	0.35	1100 (459-1501)
		Latitude	0.03	-0	0.07	1100 (777-1488)
	Random (variance)	Phylogeny	0	0	0.07	1100 (786-1393)
		Species	0	0	0.03	1100 (861-1412)
		Breeding site	0	0	0	1100 (826-1860)
		Residual	0.03	0.03	0.04	1100 (948-2039)
	Page's λ		0.74	0.02	1	1100 (843-1471)
Deviations from 24-h	Fixed	Intercept (non-tidal)	0.02	-0.04	0.09	1100 (851-1742)
		Life history (tidal)	-0.02	-0.1	0.04	1100 (702-2257)
	Random (variance)	Phylogeny	0	0	0.01	1100 (806-1692)
		Species	0	0	0	1100 (692-1601)
		Breeding site	0	0	0.01	1100 (656-1490)
		Residual	0.07	0.06	0.08	1100 (760-1563)
	Page's λ		0.77	0.01	1	1100 (864-1451)

564
565
566
567
568
569

The posterior estimates (modes) of the effect sizes with the highest posterior density intervals (95% CI) and the median and range of the effective sample sizes (N (range)) come from the joint posterior distribution of 100 separate runs each with one of the 100 separate phylogenetic trees from <http://birdtree.org>. $N_{\text{bout}} = 729$ nests from 91 populations belonging to 32 species. For models on light entrainable rhythm, absolute deviations and deviations from 24-h: $N = 584$ nests from 88 populations belonging to 30 species. Latitude (in bout model: absolute value), wing length (ln-transformed), and escape distance (ln-transformed) were mean-centred and scaled (divided by SD). The estimates for the light- entrainable rhythm are on a binomial scale. For procedures and specifications related to phylogenetic Bayesian mixed models see Methods.

570 **Extended Data Table 4 | Incubation monitoring methods and systems.**

Method	Model	Company	www	Year	Sampling interval (min)	N populations	N nests	N both parents tagged
RFID + tag embedded in the flag ^{32,33}	tag: 9.0 × 2.1 mm, 0.087 g, 134.2 kHz	Biomark	http://www.biomark.com/	2011-2014	0.08	19	200	157
	RFID: custom made	Calima Engineering, & Max Planck Institute for Ornithology	http://www.calima.de http://www.orn.mpg.de/en					
	temperature probe: TinyTag Talk PB-5005-0M6 (ø 2.5 mm) with Talk 2 logger TK-4023 or	Gemini Data Loggers Ltd	www.tinytag.info				197	
	MSR® (12 × 6 mm) with MSR® 145 logger	MSR® Electronics GmbH	http://www.msr.ch/					
	tag: 11.5 × 2.12 mm, 0.1 g, 125 kHz	Trovan®	http://www.trovan.com	2007	5	1	3	3
	RFID: custom made	EID Aalten BV	http://www.dorset.nu/en/					
RFID + tag glued to the tail ³⁴	temperature probe: TinyTag Talk PB-5005-0M6 (ø 2.5 mm) with Talk 2 logger TK-4023	Gemini Data Loggers Ltd	www.tinytag.info					
	tag: 11.5 × 2.12 mm, 0.09g, 128 kHz	Trovan®	http://www.trovan.com	2005-2008	1.7-5.5	1	34	34
	RFID: LID650 with extended memory, driver TM613, Antenna ANT614							
	tag: TIRIS 12 × 6 × 3 mm, 0.4 g, 134.2 kHz	Texas Instruments	http://www.ti.com	1997	1/3	1	6	6
	RFID: TIRIS Micro-reader							
Light logger ^{35,36,70}	TIRIS tag: 12 × 6 × 3 mm, 0.4 g, 134.2 kHz	Texas Instruments	http://www.ti.com	2006	1/3	1	18	18
	RFID: custom made	Francis Scientific Instruments Limited	https://www.duedil.com/comp-any/01964877/francis-scientific-instruments-limited					
	MK10, 12, 14 18, 20: 0.7-1.4g, light scale 0-64 (64 ≥ 100 lux)	British Antarctic Survey, currently Biotrack	http://www.biomark.com/	2008-2013	2-10	48	261*	55
	Intigeo W65A9RK: 0.7g, absolute light levels (lux)	Migrate Technology Ltd	http://www.migratetech.co.uk	2012-2014	5-10	23	136*	11
GPS-tracker ³⁷	UvA-BITS 4C: 62 × 30 × 12, 14 g	University of Amsterdam	http://www.uva-bits.nl/	2010	10-30	1	8	8
	UvA-BITS 2CDse: 52 × 22 × 9, 7.5 g			2014	20	1	1**	0
Radio-transmitter ²⁶	iTag: 26 × 15 × 9 mm, 4 g	e-obs GmbH	http://www.e-obs.de/	2009	0.07	2	3	3
Video ⁷¹⁻⁷⁴				2005-2006	1/3	1	28	-
				2005-2011	constant	3	25	-
Observations				1994	30	1	5	-
				2011	constant	1	3	-

571 For details about methods used in each populations, see Supplementary Data⁵⁴.
572 *At one nest a bird with MK logger was recaptured and the logger exchanged for Intigeo logger. This nest appears in N for both logger types.
573 **Simultaneously equipped with light logger (Intigeo). This nest appears in N for both GPS-tracer and Intigeo.